

University of Groningen

Reductie en ozonisatie van acetyleenethers

Wisaksono, Wahjoudi Wiratmoko

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1961

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wisaksono, W. W. (1961). *Reductie en ozonisatie van acetyleenethers*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SUMMARY.

In Chapter I we described the reactions of dialkylaluminiumhydrides HAL(alk)_2 with ethoxyethynylcarbinols $\text{R}-\underset{\text{R}'}{\text{C}}(\text{OH})-\text{C}\equiv$

COC_2H_5 . Several products could be isolated depending on the temperature during the hydrolysis of the initial aluminium-containing addition product. R'



At -5°C the diene ethers ($\text{R}-\underset{\text{R}'}{\text{C}}=\text{C}=\text{CHOC}_2\text{H}_5$) could be obtained together with the ethoxyvinylcarbinols. ($\text{R}-\underset{\text{R}'}{\text{C}}(\text{OH})-\text{CH}=\text{CHOC}_2\text{H}_5$).

At $+20^\circ\text{C}$ a mixture of the cumulated and conjugated diene ethers is formed. These mixtures, which also contain unidentified compounds are difficult to separate.

Only in one instance were we able to obtain a pure conjugated diene ether ($\text{CH}_2=\underset{\text{CH}_3}{\text{C}}-\text{CH}=\text{CHOC}_2\text{H}_5$).

By acid hydrolysis the ethoxyvinylcarbinols as well as the cumulated and conjugated diene ethers are converted into α, β -unsaturated aldehydes.

The reduction of ethoxyethynylcarbinols with dialkylaluminiumhydrides could therefore be used in the synthesis of non-aromatic α, β -unsaturated aldehydes.



Chapter II deals with the reactions of ozone with unsaturated ethers (vinyl ethers, acetylenic ethers, cumulated diene ethers, ethoxyethynylcarbinols).

It was possible to estimate with this reagent the concentration of the cumulated diene ethers in solution.

On ozonolysis the acetylenic ethers yielded α -keto-esters. This synthesis has some preparative value. The β -hydroxy- α -keto esters, which are not easily accessible, can thus be prepared from the ethoxyethynylcarbinols.